

EDITORIAL

Burning Questions

Burning questions: How do soil microbes shape ecosystem biogeochemistry in the context of global change?

Earth system models disagree on the future of global soil carbon C, with projections ranging from losses of 70 Pg to gains of 250 Pg by the year 2100 (Todd-Brown et al., 2014). This disagreement is due, in part, to the high degree of uncertainty in the mechanisms that control soil C (Sulman et al., 2018), the largest dynamic pool of organic C on Earth. It remains unclear how climate change, rising atmospheric CO₂ levels, and other environmental changes will combine to affect global soil C. As key engineers of biogeochemical cycling, microbes play a crucial role in soil C responses to these environmental changes. Yet, the path toward representing microbes in soil models remains uncertain. Therefore, we address four burning questions relevant for modelling microbial ecological, evolutionary, and biogeochemical processes in soils: (1) What is the role of soil microbes in biogeochemical cycling? (2) Does microbial community composition matter for soil carbon cycling? (3) Which ecological and evolutionary processes contribute to functional changes? (4) How should we model microbial eco-evolutionary mechanisms in soil C models?

QUESTION: DO MICROBES MATTER FOR SOIL BIOGEOCHEMICAL CYCLING?

Answer: Yes

As the key drivers of biogeochemical cycles, microbes are the ‘engines of life on Earth’ (Falkowski et al., 2008). Soil microbes decompose soil organic matter (SOM), with respiration responsible for over 50% of soil CO₂ emissions (Jia et al., 2016). Priming effects, whereby adding fresh carbon to soil results in the respiration of existing carbon, are impossible to explain without invoking microbial mechanisms (Kuzyakov, 2010). In addition to their role in soil carbon turnover, microbes also promote soil carbon storage. The byproducts of microbial decomposition have a high affinity for minerals, which helps to form stable soil C (Miltner et al., 2012). Consequently, soil C storage by microbes is a prospective mechanism to mitigate the effects of anthropogenic global change.

Knowledge Gaps: It is clear that microbial biomass matters for biogeochemistry, but the magnitude of

impact depends on both abiotic and biotic factors. While microbial biomass is a primary driver of decomposition in litter and the rhizosphere, other factors may dominate soil C dynamics in mineral soils, where access to substrate may limit decomposition (Schimel & Schaeffer, 2012). The identity of microbes, their traits, and their relative abundances within the community (i.e., composition) could also be important. Wieder and collaborators (Wieder et al., 2013) revealed the extreme sensitivity of soil C predictions to microbial parameters that can reflect both historical selection (phylogeny) and contemporary selection (demography) by new climatic conditions (Hanson et al., 2012; Martiny et al., 2006). Given that environmental selection varies between locations and over time, microbial parameters should reflect spatiotemporal variation. An early attempt to integrate spatio-temporal variation in microbial composition demonstrated that projections of C loss by 2100 nearly doubled (Abs et al., 2022), highlighting the need to better understand what environmental factors determine microbial properties, especially within a community context.

DOES COMMUNITY COMPOSITION MATTER FOR SOIL CARBON CYCLING?

Answer: Yes, in some cases

There are now many studies showing that biogeochemical functions, including carbon cycling, depend on microbial community composition. Based on a reciprocal transplant along a climate gradient, Glassman et al. found that rates of litter decomposition varied across different microbial communities even under the same climate and substrate conditions (Glassman et al., 2018). In laboratory microcosms, cumulative respiration of plant litter depended on microbial community composition, with the greatest respiration rates observed when microbial communities decomposed their native litter (Strickland et al., 2009). This result, and other examples of so-called ‘home-field advantage’, provide compelling evidence that microbial composition can influence soil carbon cycling (Bradford et al., 2017). Over long timescales (e.g., centuries),

processes of soil C stabilization may also be controlled by specific microbes (Schimel & Schaeffer, 2012).

On the other hand, there are conditions under which microbial community composition may have little effect on soil biogeochemistry. Some researchers initially assumed that high microbial diversity equated to high functional redundancy among community members (Lawton & Brown, 1994). Over the years, this idea has been tested, refined, and informed by trait-based theory (Allison & Martiny, 2008; Martiny et al., 2015). For example, many microbes can be considered redundant for 'broad' functions, such as respiration (Louca et al., 2018; Schimel & Schaeffer, 2012) but possibly not for 'narrow' functions, such as methane production (McCalley et al., 2014). Microbial community composition may determine C flows in litter and in the rhizosphere, but not always in mineral soil where the limiting factor is access to substrate (Dungait et al., 2012; Schimel & Schaeffer, 2012).

Knowledge Gaps: There is still work to be done to understand when and how microbial diversity is relevant for predicting biogeochemical processes. With high-throughput sequencing, characterizing who is present within a given soil microbial community is readily possible across large scales (Earth Microbiome Project). Yet, functional metrics are still often aggregated at the community level, limiting our ability to link functional observations directly to taxonomic members and mechanistically link changes in abundance to function. For example, broad genetic characterizations of microbial communities (e.g., 16 S rRNA gene amplicons) mask a high degree of variation in ecologically relevant traits among closely related strains, such as the temperature sensitivity of carbohydrate degradation (Chase et al., 2017; Johnson et al., 2006). Ultimately, there is a pressing need to understand which functions are conserved at which taxonomic depth to better understand microbial genome-to-function relationships. By elucidating the degree of trait variation among co-occurring soil microbes, and the influence of ecological processes, we can better apply trait-based frameworks to understand the impact of both taxonomic and functional variation within soil microbial communities (Malik et al., 2020).

WHAT ECOLOGICAL AND EVOLUTIONARY PROCESSES DRIVE FUNCTIONAL CHANGES?

Answer: Environmental selection, dispersal, phenotypic plasticity, mutation, horizontal gene transfer, and stochasticity may all contribute to functional changes

Ecological processes are largely associated with demography (shifts in abundance), while evolutionary processes can introduce entirely new alleles and traits into communities (Figure 1) (Chase et al., 2018). Field

experiments demonstrate that microbial community composition can respond rapidly to environmental change (Matulich et al., 2015). Concurrently, recent evidence suggests that intra-species trait variation also influences compositional changes on similar timescales (Chase et al., 2021; Garud & Pollard, 2020). Disentangling these processes provides essential information on the different mechanisms facilitating a microbiome's response to climate change.

Both taxonomic and functional changes are driven by deterministic (e.g., trait selection) and stochastic processes (Hanson et al., 2012). Deterministic processes combine more or less random sources of variation (physiological plasticity, dispersal, mutation, horizontal gene transfer) with environmental selection (sorting of species or alleles). Dispersal can stabilize or shift microbiomes through the immigration of novel and extant taxa. Stochastic processes, such as ecological and genetic drift, are non-selective and have been shown to play a major role in community assembly and functioning (Albright et al., 2018, 2019). Ultimately, a microbiome's response to environmental change involves a continuum of these processes ranging from broad taxonomic shifts to the emergence of de novo mutations (Chase et al., 2021).

Knowledge Gaps: While it is becoming increasingly clear that microbiomes respond to changing environments through ecological and evolutionary mechanisms, the timescales and relative contributions of each process remain unclear. For instance, we have a limited understanding of where microbial migrants come from (vegetation, soil, air) and whether dispersal should be viewed as an active or passive mechanism of a microbiome (Walters & Martiny, 2020). In a 30-year long climate manipulation experiment, (Melillo et al., 2017) found distinct phases of microbial community response to long-term warming. The fastest responses may result from changes in demography, whereas evolutionary responses may play out over years to decades. In addition, given that past evolutionary divergence can frequently impact contemporary ecological patterns (Martiny et al., 2017), we need to assess when and where the outcomes of evolution affect ecological processes.

Although long-term in vitro microbial evolution experiments have shown strong evidence for adaptation (Lenski, 2017; Rainey & Travisano, 1998; Rodríguez-Verdugo, 2021; Travisano & Lenski, 1996), it is unclear if these observations extend to natural systems (Koskella & Vos, 2015). In the few examples investigating the impact of evolution within natural communities, it appears that slow generation times (Caro et al., 2022) combined with high spatial heterogeneity may limit detection of genome evolution and result in different evolutionary dynamics than those observed in laboratory environments (Chase et al., 2021). Resolving the feedbacks between ecological and evolutionary processes will be essential for improving model predictions of




FIGURE 1 Conceptual diagram of microbial eco-evolutionary feedbacks to global change (here warming) and their implications for global soil C stocks.

biogeochemical functions, and might contradict the current belief that biodiversity inhibits evolutionary responses to changing environments (de Mazancourt et al., 2008; Johansson, 2008; Loeuille & Leibold, 2008).

HOW SHOULD WE MODEL MICROBIAL ECO-EVOLUTIONARY MECHANISMS IN SOIL C MODELS?

Answer: We need to represent observed mechanisms in models and develop approaches for scaling them up

Thus far, microbial updates to soil biogeochemical models have involved very coarse representations of eco-evolutionary mechanisms. Several models have integrated the acceleration of microbial enzyme kinetics (degradation, uptake) with warming (biochemical response) (Sulman et al., 2014; Wieder et al., 2013; Wieder et al., 2014, 2015). A few models have also added a linear decrease in microbial carbon use efficiency (CUE) with warming (Li et al., 2014), which has been observed in short-term warming experiments and is the result of higher cellular maintenance needs (physiological response) (Geyer et al., 2019; Giorgio, &

del, and Jonathan J. Cole., 1998; Manzoni et al., 2018). Wieder et al.'s model assumed that if microbes can adapt to warming, they will reduce or even cancel that loss in CUE (Wieder et al., 2013). They found that decreasing CUE (short-term physiological response) led to a slight soil C gain (5 Pg) by 2100, while constant CUE (long-term adaptation) led to a large soil C loss (300 Pg). However, there is no proof that evolutionary pressure will keep CUE constant.

More mechanistic modelling approaches are starting to emerge. Abs et al. proposed a mathematical method based on game theory to predict the microbial community eco-evolutionary response to warming (Abs et al., 2022). This method accounts for progressive community trait change modifying the ecological environment (resources, competition), which initiates an eco-evolutionary feedback to modify selection of community traits. The model is agnostic to the ecological versus evolutionary mechanisms contributing to the trait change. The model predicted that microbes' C allocation to the production of C-targeting enzymes should non-linearly increase with warming, leading to greater soil C loss than in the non-adaptive model and with losses concentrated in cold regions.

Knowledge Gaps: Although there has been some recent progress in modelling microbial eco-evolutionary

mechanisms, there are still many open questions. Do different mechanisms lead to different community functions, or do they converge functionally? Does it matter whether dispersal or mutation introduces new alleles, or can they be modelled as a combined process? Should the emergence of new variants be modelled as a constant rate or a variable dependent on each process? Empirical evidence suggests a complicated interaction between ecological and evolutionary processes (Chase et al., 2021). Consequently, eco-evolutionary theoretical models that predict the relative contribution of each process (e.g., demography vs. mutations) will need to account for environmental selective pressures across different timescales. For instance, we predict that (1) new functions arising from de novo mutations might dominate in highly fluctuating, dispersal-limited environments; (2) dispersal could facilitate rapid functional turnover from regional species or population pools; and (3) demographic shifts could be more pronounced in isolated, functionally diverse communities and slow-oscillating environments (Loeuille & Leibold, 2008).

CONCLUSION

We know that microbial communities respond to climate change through processes occurring at many spatiotemporal scales, ranging from physiological acclimation to entire community compositional shifts. Yet, it remains unclear whether incorporating these processes into ecosystem models will reduce the current uncertainty in soil C predictions. The paradigm shift from modelling soil microbial communities as a 'black box' to appreciating their role in soil formation has certainly propelled soil C modelling into a new, integrative frontier. With more local-scale empirical studies disentangling ecological and evolutionary processes, we can better understand whether and how adaptive processes influence predictions of microbial functioning. The integration of realistic soil spatial structure (pores, aggregates, rhizosphere vs. bulk soil) is another necessary step toward reducing uncertainty in soil biogeochemical models. By taking these steps, we can build better mechanistic models of microbial community dynamics. With these models in hand, we can predict the impact of microbes on soil C balance with applications for future climate prediction and greenhouse gas mitigation.

AUTHOR CONTRIBUTIONS

Elsa Abs: Conceptualization (lead); investigation (equal); supervision (equal); validation (equal); writing – original draft (lead); writing – review & editing (lead). **Alexander Chase:** Conceptualization (equal); investigation (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Steven**

D. Allison: Validation (equal); writing – review and editing (equal).

ACKNOWLEDGEMENTS

We thank Mallory Choudoir, Ashish Malik and Adriana Romero-Olivares for their friendly review, whose comments really improved this article. Funding was provided to Elsa Abs by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 891576; to Steven D. Allison by the US National Science Foundation (DEB-2016482) and the US Department of Energy, Office of Science, BER (SC-0020382).

CONFLICT OF INTEREST

The authors declare no conflict of interest.


FUNDING INFORMATION

H2020 Marie Skłodowska-Curie Actions, Grant/Award Number: 891576; U.S. Department of Energy, Grant/Award Number: SC-0020382; European Union; Horizon 2020; National Science Foundation, Grant/Award Number: DEB-2016482; Department of Energy; Office of Science

DATA AVAILABILITY STATEMENT

No data, no code was used to make this manuscript.

Elsa Abs¹ 

Alexander B. Chase² 

Steven D. Allison^{1,3} 

- ¹Department of Ecology and Evolutionary Biology, University of California, Irvine, California, USA
²Department of Earth Sciences, Southern Methodist University, Dallas, Texas, USA
³Department of Earth System Science, University of California, Irvine, California, USA

Correspondence

Elsa Abs, Department of Ecology and Evolutionary Biology, University of California, Irvine, CA, USA.
 Email: eabs@uci.edu

ORCID

Elsa Abs  <https://orcid.org/0000-0001-9501-1412>

Alexander B. Chase  <https://orcid.org/0000-0003-1984-6279>

Steven D. Allison  <https://orcid.org/0000-0003-4629-7842>

REFERENCES

- Abs, E., Saleska, S. & Ferriere, R. (2022) Microbial eco-evolutionary responses amplify global soil carbon loss with climate warming. <https://doi.org/10.21203/rs.3.rs-1984500/v1>

- Albright, M.B.N. & Jennifer, B.H.M. (2018) Dispersal alters bacterial diversity and composition in a natural community. *The ISME Journal*, 12(1), 296–299.
- Albright, M.B.N., Alexander B.C. & Jennifer, B.H.M. (2019) Experimental evidence that stochasticity contributes to bacterial composition and functioning in a decomposer community. *mBio*, 10(2). Available from: <https://doi.org/10.1128/mBio.00568-19>
- Allison, S.D. & Martiny, J.B.H. (2008) Resistance, resilience, and redundancy in microbial communities. *Proceedings of the National Academy of Sciences*, 105(1), 11512–11519.
- Bradford, M.A., Veen, G.F., Bonis, A., Bradford, E.M., Classen, A.T., Hans, J. et al. (2017) A test of the hierarchical model of litter decomposition. *Nature Ecology & Evolution*, 1, 1836–1845. Available from: <https://doi.org/10.1038/s41559-017-0367-4>
- Caro, T.A., McFarlin, J., Jech, S., Fierer, N. & Kopf, S. (2022) Hydrogen stable isotope probing of lipids demonstrates slow rates of microbial growth in soil. *bioRxiv*. Available from: <https://doi.org/10.1101/2022.07.11.499392>
- Chase, A.B., Gomez-Lunar, Z., Lopez, A.E., Li, J., Allison, S.D., Martiny, A.C. et al. (2018) Emergence of soil bacterial ecotypes along a climate gradient. *Environmental Microbiology*, 20(11), 4112–4126.
- Chase, A.B., Karaoz, U., Brodie, E.L., Gomez-Lunar, Z., Martiny, A.C. & Martiny, J.B.H. (2017) Microdiversity of an abundant terrestrial bacterium encompasses extensive variation in ecologically relevant traits. *mBio*, 8(6), e01809-17. Available from: <https://doi.org/10.1128/mBio.01809-17>
- Chase, Alexander B., Claudia Weihe, and Jennifer B. H. Martiny. 2021. “Adaptive differentiation and rapid evolution of a soil bacterium along a climate gradient. *Proceedings of the National Academy of Sciences of the United States of America*. 118(18), e2101254118. <https://doi.org/10.1073/pnas.2101254118>.
- de Mazancourt, C., Johnson, E. & Barraclough, T.G. (2008) Biodiversity inhibits Species' evolutionary responses to changing environments. *Ecology Letters*, 11(4), 380–388.
- Del Giorgio, P.A. & Jonathan J. Cole. (1998) Bacterial growth efficiency in natural aquatic systems. *Annual Review of Ecology and Systematics*, 29(1), 503–541.
- Dungait, J.A.J., Hopkins, D.W., Gregory, A.S. & Whitmore, A.P. (2012) Soil organic matter turnover is governed by accessibility not recalcitrance. *Global Change Biology*, 18(6), 1781–1796.
- Falkowski, P.G., Fenchel, T. & Delong, E.F. (2008) The microbial engines that drive Earth's biogeochemical cycles. *Science*, 320(5879), 1034–1039.
- Garud, N.R. & Pollard, K.S. (2020) Population genetics in the human microbiome. *Trends in Genetics*, 36(1), 53–67.
- Geyer, K.M., Dijkstra, P., Sinsabaugh, R. & Frey, S.D. (2019) Clarifying the interpretation of carbon use efficiency in soil through methods comparison. *Soil Biology & Biochemistry*, 128 (January), 79–88.
- Glassman, S.I., Weihe, C., Li, J., Albright, M.B.N., Looby, C.I., Martiny, A.C. et al. (2018) Decomposition responses to climate depend on microbial community composition. *Proceedings of the National Academy of Sciences of the United States of America*, 115(47), 11994–11999.
- Hanson, C.A., Fuhrman, J.A., Claire Horner-Devine, M. & Martiny, J. B.H. (2012) Beyond biogeographic patterns: processes shaping the microbial landscape. *Nature Reviews. Microbiology*, 10(7), 497–506.
- Jia, S., Zhang, X., Chen, X., McLaughlin, N.B., Zhang, S., Wei, S. et al. (2016) Long-term conservation tillage influences the soil microbial community and its contribution to soil CO₂ emissions in a Mollisol in Northeast China. *Journal of Soils and Sediments*, 16(1), 1–12.
- Johansson, J. (2008) Evolutionary responses to environmental changes: how does competition affect adaptation? *Evolution; International Journal of Organic Evolution*, 62(2), 421–435.
- Johnson, Z.I., Zinser, E.R., Coe, A., McNulty, N.P., Malcolm, E., Woodward, S. et al. (2006) Niche partitioning among *Prochlorococcus* ecotypes along ocean-scale environmental gradients. *Science*, 311(5768), 1737–1740.
- Koskella, B. & Vos, M. (2015) Adaptation in natural microbial populations. *Annual Review of Ecology, Evolution, and Systematics*, 46(1), 503–522.
- Kuzyakov, Y. (2010) Priming effects: interactions between living and dead organic matter. *Soil Biology & Biochemistry*, 42(9), 1363–1371.
- Lawton, J.H. & Brown, V.K. (1994) Redundancy in ecosystems. In: Schulze, E.-D. & Mooney, H.A. (Eds.) *Biodiversity and Ecosystem Function*. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 255–270.
- Lenski, R.E. (2017) Convergence and divergence in a long-term experiment with bacteria. *The American Naturalist*, 190(S1), S57–S68.
- Li, J., Wang, G., Allison, S.D., Mayes, M.A. & Luo, Y. (2014) Soil carbon sensitivity to temperature and carbon use efficiency compared across microbial-ecosystem models of varying complexity. *Biogeochemistry*, 119(1), 67–84.
- Loeuille, N. & Leibold, M.A. (2008) Evolution in Metacommunities: on the relative importance of species sorting and monopolization in structuring communities. *The American Naturalist*, 171(6), 788–799.
- Louca, S., Polz, M.F., Mazel, F., Albright, M.B.N., Huber, J.A., O'Connor, M.I. et al. (2018) Function and functional redundancy in microbial systems. *Nature Ecology & Evolution*, 2(6), 936–943.
- Malik, A.A., Martiny, J.B.H., Brodie, E.L., Martiny, A.C., Treseder, K. K. & Allison, S.D. (2020) Defining trait-based microbial strategies with consequences for soil carbon cycling under climate change. *The ISME Journal*, 14(1), 1–9.
- Manzoni, S., Čapek, P., Porada, P., Thurner, M., Winterdahl, M., Beer, C. et al. (2018) Reviews and syntheses: carbon use efficiency from organisms to ecosystems—definitions, theories, and empirical evidence. *Biogeosciences*, 15(19), 5929–5949. Available from: <https://doi.org/10.5194/bg-15-5929-2018>
- Martiny, J.B., Hughes, J.B., Martiny, H., Bohannan, B.J.M., Brown, J. H., Colwell, R.K. et al. (2006) Microbial biogeography: putting microorganisms on the map. *Nature Reviews Microbiology*, 4, 102–112. Available from: <https://doi.org/10.1038/nrmicro1341>
- Martiny, J.B.H., Jones, S.E., Lennon, J.T. & Martiny, A.C. (2015) Microbiomes in light of traits: a phylogenetic perspective. *Science*, 350, aac9323. Available from: <https://doi.org/10.1126/science.aac9323>
- Martiny, J.B., Martiny, A.C., Weihe, C., Ying, L., Berlemont, R., Brodie, E.L. et al. (2017) Microbial legacies alter decomposition in response to simulated global change. *The ISME Journal*, 11(2), 490–499.
- Matulich, K.L., Weihe, C., Allison, S.D., Amend, A.S., Berlemont, R., Goulden, M.L. et al. (2015) Temporal variation overshadows the response of leaf litter microbial communities to simulated global change. *The ISME Journal*, 9(11), 2477–2489.
- McCalley, C.K., Woodcroft, B.J., Hodgkins, S.B., Wehr, R.A., Kim, E.-H., Mondav, R. et al. (2014) Methane dynamics regulated by microbial community response to permafrost thaw. *Nature*, 514(7523), 478–481.
- Melillo, J.M., Frey, S.D., DeAngelis, K.M., Werner, W.J., Bernard, M. J., Bowles, F.P. et al. (2017) Long-term pattern and magnitude of soil carbon feedback to the climate system in a warming world. *Science*, 358(6359), 101–105.
- Miltner, A., Bombach, P., Schmidt-Brücken, B. & Kästner, M. (2012) SOM genesis: microbial biomass as a significant source. *Biogeochemistry*, 111(1), 41–55.
- Rainey, P.B. & Travisano, M. (1998) Adaptive radiation in a heterogeneous environment. *Nature*, 394(6688), 69–72.

- Rodríguez-Verdugo, A. (2021) Evolving interactions and emergent functions in microbial consortia. *mSystems*, 6(4), e0077421.
- Schimel, J.P. & Schaeffer, S.M. (2012) Microbial control over carbon cycling in soil. *Frontiers in Microbiology*, 3(September), 348.
- Strickland, M.S., Lauber, C., Fierer, N. & Bradford, M.A. (2009) Testing the functional significance of microbial community composition. *Ecology*, 90(2), 441–451.
- Sulman, B.N., Moore, J.A.M., Abramoff, R., Averill, C., Kivlin, S., Georgiou, K. et al. (2018) Multiple models and experiments underscore large uncertainty in soil carbon dynamics. *Biogeochemistry*, 141(2), 109–123.
- Sulman, B.N., Phillips, R.P., Christopher Oishi, A., Shevliakova, E. & Pacala, S.W. (2014) Microbe-driven turnover offsets mineral-mediated storage of soil carbon under elevated CO₂. *Nature Climate Change*, 4(12), 1099–1102. Available from: <https://doi.org/10.1038/nclimate2436>
- Todd-Brown, K.E.O., Randerson, J.T., Hopkins, F., Arora, V., Hajima, T., Jones, C. et al. (2014) Changes in soil organic carbon storage predicted by earth system models during the 21st century. *Biogeosciences*, 10(12), 18969–19004. Available from: <https://doi.org/10.5194/bg-11-2341-2014>
- Travisano, M. & Lenski, R.E. (1996) Long-term experimental evolution in *Escherichia coli*. IV. Targets of selection and the specificity of adaptation. *Genetics*, 143(1), 15–26.
- Walters, K.E. & Jennifer, B.H.M. (2020) Alpha-, Beta-, and Gamma-Diversity of Bacteria Varies across Habitats. *PLoS One*, 15(9), e0233872.
- Wieder, W.R., Grandy, A.S., Kallenbach, C.M. & Bonan, G.B. (2014) Integrating microbial physiology and physio-chemical principles in soils with the microbial-mineral carbon stabilization (MIMICS) model. *Biogeosciences*, 11(14), 3899–3917. Available from: <https://doi.org/10.5194/bg-11-3899-2014>
- Wieder, W.R., Grandy, A.S., Kallenbach, C.M., Taylor, P.G. & Bonan, G.B. (2015) Representing life in the earth system with soil microbial functional traits in the MIMICS model. *Geoscientific Model Development Discussions*, 8(2), 2011–2052.
- Wieder, W.R., Bonan, G.B. & Allison, S.D. (2013) Global soil carbon projections are improved by modelling microbial processes. *Nature Climate Change*, 3(10), 909–912.